Antibiogram Reports



- ✓ Simple, secure downloading ✓ Exceeds standards and guidelines
- ✓ Includes summary resistance patterns

YOUR ANTIBIOGRAM REPORT

Assists you to comply with items 3.2 and 3.14 of the National Safety and Quality Health Service Standard 3.

Shows the resistance patterns of your organisms for the previous year.

Useful to review your empiric antibiotic guidelines.

According to national and international guidelines, only the first isolate of an organism recovered from each patient during the year is used to calculate the percentage of susceptibility for each given species.

This is done to prevent the over reporting of resistance.

You will also have access to the raw data that is used to calculate the percentages of sensitivity.

Methods used are based upon the recommendations of the Clinical and Laboratory Standards Institute (CLSI) and the Australian Commission on Safety and Quality in Health Care.





Antibiogram Reports



>>> CONTINUED FROM OVERLEAF

YOUR ANTIBIOGRAM REPORT PRESENTS:

- Blood culture data full antibiogram (top 10 organisms isolated for your institution)
- Urine culture data full antibiogram (top 5 organisms isolated for your institution)
- Collated non-urine culture data full antibiogram (top 10 isolated for your institution)
- 15 specific organisms with full antibiogram data from your institution
- **Separate raw patient data** provided (includes multiple isolates from the same patient)

PROCESS

We require your Medical Director or CEO to inform us that you require access to this annual report. We will create a secure user name and password for you to access the reports. Reports can be received by different recipients to the available 'Infection Control Reports'.

| 4 | Α | В | С | D | E | F | G | Н | -1 | J | K | L | M | N | 0 | Р | Q | R | S | Т | U | V | W |
|----------|------|--|-----------------------|----------------|-------------|-----------|------------|-------------|-------------|---------------|----------------|-------------|------------|------------|------------|---------|----------|----------------|----------|-----------|------------|--------------|--------------|
| 1 | Anti | biogram 2012 (Jan-Dec) for 'XY | Z' Hospital : | Blood (| Culture | es Iso | ates | | | | | | | | | | | | | | | | |
| 2 | NOS | Organism | Total nos of isolates | Data | Amoxycillin | Augmentin | Cephalexin | Cephalothin | Clindamycin | Cotrimoxazole | Flucioxacillin | FusidicAcid | Gentamicin | Penicillin | Rifampicin | Tazodin | Timentin | Trimethoprim | Amikadn* | Cefepime* | Cefoxitin* | Ceftazidime* | Ceftriaxone* |
| 3 | 1 | Escherichia coli | 130 | %Sens | 48.1 | 79.8 | 54.8 | 46.2 | | 76.9 | | | 96.9 | | | 94.6 | 82 | 100 | 100 | 97.3 | 96.7 | 96.3 | 96. |
| 4 | | | | Total | 129 | 129 | 62 | 13 | | 130 | | | 130 | | | 92 | 128 | 3 | 2 | 112 | 92 | 108 | 13 |
| 5 | 2 | Coagulase negative Staphylococcus | | %Sens | | | | | 68.8 | 70 | 47.7 | 75 | 85.7 | 15.3 | 90 | | | | | | | | |
| 6 | | | | Total | | | | | 109 | 20 | 111 | 20 | 21 | 111 | 20 | | | - | | | | | |
| 7 | 3 | Staphylococcus aureus - ALL | | %Sens | | | | | 75.3 | 94.9 | 82.2 | 100 | 94.9 | 12.3 | | | | | | | | | |
| 8 | | | | Total | | | | | 73 | 59 | 73 | 60 | 59 | 73 | 60 | | | | | | | | |
| 9 | 4 | Staphylococcus aureus - MSSA | | %Sens | | | | | 88.3 | 100 | 100 | 100 | 100 | 15 | | | | | | | | | |
| 10 | | | | Total | | | | | 60 | 49 | 60 | 49 | 49 | 60 | 49 | | | - | | | | | |
| 11 | 5 | Streptococcus pneumoniae | | %Sens | | | | | 80 | | | | | 94.4 | | | | | | | | | 10 |
| 12 | _ | | | Total | 400 | | | | 20 | | | | | 18 | | | | | | | | | 2 |
| 13 | 6 | Enterococcus faecalis | | %Sens | 100 | | | | 0 | | | | | 83.3 | | | | | | | | | |
| 14 15 | _ | Standard Standard Standard | | Total | 16 | | | | 5 | 70 | | 400 | 70 | 6 | 400 | | | | | | | | 1 |
| 16 | 7 | Staphylococcus aureus - MRSA | | %Sens Total | | | 0 12 | | 15.4 | 70 | 0 | 100 11 | 70 10 | 0 13 | 100 11 | | | | | | | | |
| 17 | 8 | Staph. epidermidis | | %Sens | | | 12 | | 13 50 | 10 60 | 13 25 | 70 | 80 | 8.3 | | | | | | | | | |
| 18 | 8 | Stapn. epidermidis | | %Sens Total | | | | | 12 | 10 | 12 | 10 | 10 | 12 | 100 | | | | | | | | |
| 19 | 9 | Beta haem. strep. (Group B) | | %Sens | | | | | 70 | 10 | 12 | 10 | 10 | 100 | 10 | | | | | | | | 10 |
| 20 | 9 | beta flaem. strep. (Group b) | | Total | | | | | 10 | | | | | 100 | | | | : | | | | | 1 |
| 21 | 10 | Klebsiella pneumoniae | | %Sens | 0 | 100 | 100 | | 10 | 100 | | | 100 | 10 | | 100 | 100 | - i | | 100 | 100 | 100 | 10 |
| 22 | 10 | Riebsiena prieumomae | | Total | 8 | 8 | 4 | | | 8 | | | 8 | | | 5 | 8 | | | 8 | 6 | 8 | |
| 23 | | Total Positive | 380 | Total | - | | - | | | | | | | | | | | - | | | • | | |
| 24 | | % analysed of all Blood Cultures | 74.2 | | | | | | | | | | | | | | | | | | | | |
| 25 | KEY | 70 direction of the proof curtaines | 7 | | | | | | | | | | | | | | | i | | | | | -1-1-1 |
| | | of isolates sensitive: green shading | | | | | | | - | | | | | | | | | | | | | | |
| | | % of isolates sensitive: yellow shading | | | | | | | | | | | | | | | | | | | | | |
| 28 | | of isolates sensitive: red shading | | | | | | | | | | | | | | | | | | | | | |
| 29 | | tibiotics usually restricted: stippled sha | ding | | | | | | | | | | | | | | | | | | | | |
| 30 | | tibiotic not recommended to be used in | | ut specia | list advi | ce | | | | | | | | | | | | | | | | | |

Each worksheet shows resistance patterns of specific organisms in your institution.

FURTHER INFORMATION

For more information, please email: **AntibiogramReport@healius.com.au** or to discuss the Antibiogram Control Report, please contact one of our Medical Microbiologists on **(02) 9005 7606.**

